

Quantum Interference and the Trapped Bose Condensed System

Juhao Wu and A. Widom ^a

^aPhysics Department, Northeastern University, Boston MA 02115

In experiments involving Bose condensed atoms trapped in magnetic bottles, plugging the hole in the bottle potential with a LASER beam produces a new potential with two minima, and thus a condensate order parameter (i.e. wave function) with two maxima. When the trapping potential is removed and the condensate explodes away from the trap, the two wave function maxima act as two coherent sources which exhibit amplitude interference. A simplified theoretical treatment of this experimental effect is provided by considering momentum distributions.

PACS numbers: 03.75.Fi, 05.30.Jp, 32.80.Pj, 64.60.-i

The two fluid[1] model has long served as the standard picture of superfluidity in liquid He^4 . In spite of the great success of the model, Landau warned[2] that the view was merely quasi-classical. It would be desirable to have a quantum fluid mechanical model in which superposition of amplitudes on a macroscopic scale played a more central role. For the dilute Bose condensed gas[3], such coherence would mean that the Bose condensate order parameter[4] $\Psi(\mathbf{r})$ would be required to maintain many of the properties normally attributed to single particle wave functions exhibiting amplitude superposition[5].

Recent experimental probes[6], developed for observing Bose condensates[7] in atomic traps[8], have provided that which is needed to go beyond the Landau “quasi-classical” two fluid picture. Direct observation of amplitude superposition in the Bose condensate order parameter $\Psi(\mathbf{r})$ has been observed[9] in mesoscopic samples.

Consider some central features of recent experimental methods. To see what is involved, let us first contemplate a “thought experiment”. Suppose that at times $t < 0^-$, there is but one atom localized near the origin in the region of a trap potential $U(\mathbf{r})$. At time $t = 0$, the trap potential is suddenly removed, and thereafter $t > 0^+$ the atomic motion is described by the free particle Schrödinger equation $i\hbar(\partial\psi/\partial t) = -(\hbar^2/2M)\nabla^2\psi$.

From the viewpoint of quantum mechanics, the

sudden[10] removal of the trap potential yields an out going wave function described using the non-relativistic propagator[11] $G(\mathbf{r} - \mathbf{s}, t)$. The outgoing wave function is determined by

$$\psi(\mathbf{r}, t) = \int G(\mathbf{r} - \mathbf{s}, t)\psi(\mathbf{s}, 0)d^3s, \quad (1)$$

where

$$G(\mathbf{r} - \mathbf{s}, t) = \left(\frac{M}{2\pi i\hbar t}\right)^{3/2} \exp\left(\frac{iM|\mathbf{r} - \mathbf{s}|^2}{2\hbar t}\right). \quad (2)$$

From the viewpoint of classical mechanics, the particle exits the trap region at a uniform velocity along the path $\mathbf{r} = \mathbf{v}t$. Both the classical (measurement) and quantum (virtual) viewpoints are resolved by considering what happens after the previously trapped particle freely moves far away from the original trapping region. Employing $|\mathbf{r}| \gg |\mathbf{s}|$ in Eqs.(1) and (2), one finds that

$$\psi(\mathbf{r}, t) \approx (M/(2\pi i\hbar t))^{3/2} e^{iS(\mathbf{r}, t)/\hbar} A(M\mathbf{r}/t), \quad (3)$$

where $S(\mathbf{r}, t) = (M|\mathbf{r}|^2/2t)$ and

$$A(\mathbf{p}) = \int e^{-i(\mathbf{p}\cdot\mathbf{s})/\hbar}\psi(\mathbf{s}, 0)d^3s. \quad (4)$$

is the momentum amplitude of the particle when it was trapped by the potential. Eq.(3) implies that

$$|\psi(\mathbf{r}, t)|^2 \approx \int |A(\mathbf{p})|^2 \delta(\mathbf{r} - (\mathbf{p}t/M)) \frac{d^3p}{(2\pi\hbar)^3}. \quad (5)$$

Thus, when a particle has escaped away from the trapping region (long after the trap potential has been removed), the probability density in space $|\psi(\mathbf{r}, t)|^2$ can be computed *by averaging the classical equation of motion* $\mathbf{r} = (\mathbf{p}t/M)$ over the momentum probability density $|A(\mathbf{p})|^2$, present when the particle was in a trapped state.

To understand recent experiments on quantum interference for trapped Bose condensed mesoscopic systems, one must replace the above single particle amplitudes with their second quantized operator counterparts

$$\hat{\psi}(\mathbf{r}) = \int e^{i(\mathbf{p} \cdot \mathbf{r})/\hbar} \hat{A}(\mathbf{p}) \frac{d^3p}{(2\pi\hbar)^3}. \quad (6)$$

The many Bosons are held in the trap for times $t < 0$ with an equilibrium momentum distribution $N(\mathbf{p}) = \langle \hat{A}^\dagger(\mathbf{p}) \hat{A}(\mathbf{p}) \rangle$.

The atomic interaction potentials have importance in assuring that the momentum distribution $N(\mathbf{p})$ (for the Bosons within the trap potential) actually correspond to thermal equilibrium. The experimentalist at time $t = 0$ *suddenly turns off the trap potential*[12] and optically monitors the particle density for times $t > 0$. The density is determined by

$$\rho(\mathbf{r}, t) = \langle \hat{\psi}^\dagger(\mathbf{r}, t) \hat{\psi}(\mathbf{r}, t) \rangle, \quad (8)$$

after the Bosons move away from the original trapping potential spatial region.

To the extent that the interaction potentials have a negligible effect on the particles after they explode away from the trapping region, the final mean density in space $\rho(\mathbf{r}, t)$ is related to the initial mean momentum distribution $N(\mathbf{p})$ via the many body version of Eq.(5); i.e.

$$\rho(\mathbf{r}, t) \approx \int N(\mathbf{p}) \delta(\mathbf{r} - (\mathbf{p}t/M)) \frac{d^3p}{(2\pi\hbar)^3}. \quad (9)$$

The *ballistic* Eq.(9) has been the central theoretical tool employed in the recent measurements of trapped Bose condensed systems. *The momentum distribution of particles within the trap is probed by the spatial distribution of particles detected away from the trap.*

Let us now review the notion of superposition of amplitudes as it becomes apparent in

the classic “two slit quantum interference experiment” [13]. Quantum particle diffraction experiments present us with a closely analogous situation regarding initial probability distributions in momentum and final probability distributions in space. In standard slit diffraction experiments, *the momentum distribution of particles just behind the slits is probed by the spatial distribution of the particles at the detectors very far behind from slits.* Once this close analogy between diffraction experiments and recent Bose condensed experiments is appreciated, it then becomes evident why these measurements have exhibited interference effects on the mesoscopic scale of the trapped Bose condensates.

Our brief review of diffraction theory from slits is as follows: (i) A particle with high velocity v is normally incident on a screen with one or more slits. The slits have a length scale $\bar{d}$ and the high incoming momentum is defined by the inequality $Mv \gg (\hbar/\bar{d})$. (ii) After a time period

$$t_s \approx (D/v) \quad (\text{slit to detector time}), \quad (10)$$

the particle hits some detector at a distance D behind the slits. (iii) Before the particle hits the slits from incidence side of the screen, the momentum parallel to the screen is zero. If the particle is squeezed through the slits to the emission side of the screen at time zero, then the quantum mechanical uncertainty principle forces the momentum amplitude

$$A(p) = \int_{\text{slits}} e^{-i(px')/\hbar} \psi(x', 0) dx'. \quad (11)$$

The momentum component p is in the x -direction, and lives in the plane of the screen with the slits. In that plane, p is normal to the long slit direction. (iv) Far behind the slits, the spatial distribution in x near the detectors is given by a one-dimensional version of Eq.(5); It is

$$|\psi(x, t_s)|^2 \approx \int |A(p)|^2 \delta(x - (pt_s/M)) \frac{dp}{(2\pi\hbar)}. \quad (12)$$

From Eqs.(10) and (12) follows the general slit diffraction formula

$$|\psi(x, D)|^2 \approx \left(\frac{Mv}{2\pi\hbar D} \right) \left| A(p = (xMv/D)) \right|^2. \quad (13)$$

For example, for diffraction through one slit having a width w , Eq.(11) reads

$$A_{(1-slit)}(p) = \frac{1}{\sqrt{w}} \int_{-w/2}^{w/2} e^{-i(px')/\hbar} dx', \quad (14)$$

so that Eq.(13) yields the one slit diffraction pattern

$$|\psi_{(1-slit)}(x)|^2 \approx \left(\frac{Mvw \sin^2(Mvwx/2\hbar D)}{2\pi\hbar D(Mvwx/2\hbar D)^2} \right). \quad (15)$$

For two slits, each of width w and separated by a distance b which obeys $b > w$, the momentum amplitudes obey

$$A_{(2-slit)}(p) = \sqrt{2} \cos\left(\frac{pb}{2\hbar}\right) A_{(1-slit)}(p), \quad (16)$$

leading to the diffraction pattern

$$|\psi_{(2-slit)}(x)|^2 \approx 2 \cos^2\left(\frac{Mvbx}{2\hbar D}\right) |\psi_{(1-slit)}(x)|^2. \quad (17)$$

Thus, one sees that the momentum distribution near the slits give rise to the diffraction pattern in space near the detectors, via Eq.(13). Let us now turn to the quantum interference diffraction pattern expected from the momentum distribution of a trapped Boson condensate.

In the zero temperature Gross[14]-Pitaevskii[15] theory for dilute Bosons in a trap potential $U(\mathbf{r})$, the variational trial ground state is taken to be “quantum coherent” in a sense that is closely analogous to coherent states in quantum optics. The trial ground state is a normalized eigenstate of the Boson field operator,

$$\hat{\psi}(\mathbf{r})|\Psi\rangle = \Psi(\mathbf{r})|\Psi\rangle. \quad (18)$$

The Bose condensation order parameter $\Psi(\mathbf{r})$ is then chosen to minimize the energy functional

$$\mathcal{E} = \langle \Psi | \hat{H} - \mu \hat{N} | \Psi \rangle \quad (19)$$

where μ is the chemical potential. Eq.(19) is evaluated using the two body scattering length a ;

$$\mathcal{E} = \int \left\{ \Psi^*(h - \mu)\Psi + (2\pi\hbar^2 a/M)|\Psi|^4 \right\} d^3r, \quad (20)$$

where

$$h = -\left(\frac{\hbar^2}{2M}\right)\nabla^2 + U(\mathbf{r}). \quad (21)$$

Minimizing Eq.(20) yields

$$\hbar\Psi(\mathbf{r}) + (4\pi\hbar^2 a|\Psi(\mathbf{r})|^2/M)\Psi(\mathbf{r}) = \mu\Psi(\mathbf{r}). \quad (22)$$

In the trapped Boson experiments, one starts the design of the trap with an inhomogeneous magnetic field so that $U_0(\mathbf{r})$ behaves as an anisotropic oscillator potential with a single minimum at the origin. Such a single minimum in $U_0(\mathbf{r})$ yields for the order parameter $\Psi_0(\mathbf{r})$ a single maximum. This single maximum in $\Psi_0(\mathbf{r})$ would behave as single source for the condensate explosion once the trap potential is removed. However, the single minimum in $U_0(\mathbf{r})$ (i.e. the single maximum in $\Psi_0(\mathbf{r})$) system is hard to build because the integer oriented spin atoms slip through a “hole” at the trapping potential minimum.

To fix this potential, some workers have plugged up the hole at the bottom of the magnetic trap with a LASER beam. The plug produces (as a side effect) an effective $U(\mathbf{r})$ with *two minima*. Thus, traps with a plug can be (and have been) designed[9] so that the order parameter $\Psi(\mathbf{r})$ has *two maxima*. In such a case, when the trap is removed and the condensate explodes away from the trap region, there will be a *double source* for the condensate Bosons. If these two sources are *coherent*, then there was in the laboratory amplitude interference yielding order parameter fringes. These fringes were known to be analogous to the interference fringes observed in conventional “two slit diffraction”. If the two sources were *incoherent*, then such interference fringes would not have occurred.

It is to be expected that Bose condensed systems (even at finite temperature) will exhibit a well defined coherent order parameter $\Psi(\mathbf{r})$ whose condensate momentum distribution

$$N_0(\mathbf{p}) = \left| \int e^{-i(\mathbf{p}\cdot\mathbf{r})/\hbar} \Psi(\mathbf{r}) d^3r \right|^2 \quad (23)$$

will properly reflect the two maxima in $\Psi(\mathbf{r})$. Such a double maxima describing two coherent sources duplicate in $\rho(\mathbf{r},t)$ the interference pattern closely analogous to two slit diffraction. After the trapping potential is removed, the condensate atom contribution to the density is given

by

$$\rho_0(\mathbf{r}, t) \approx \int N_0(\mathbf{p}) \delta(\mathbf{r} - (\mathbf{p}t/M)) \frac{d^3p}{(2\pi\hbar)^3} . \quad (24)$$

Reasoning as in Eqs.(16) and (17), and employing a separation vector $\mathbf{b}$ which describes the position of the second peak in $\Psi(\mathbf{r})$ relative to the first peak in $\Psi(\mathbf{r})$, we find that

$$N_{0(2\text{-peak})}(\mathbf{p}) \approx I(\mathbf{p}, \mathbf{b}) N_{0(1\text{-peak})}(\mathbf{p}), \quad (25)$$

where the interference function

$$I(\mathbf{p}, \mathbf{b}) = (1 + \eta \cos(\mathbf{p} \cdot \mathbf{b}/\hbar)), \quad (26)$$

and $0 < \eta < 1$ describes the “visibility” of the interference fringes when the two peaks (i.e. sources) are only partially coherent.

Finally, we note that the oscillations in momentum $\mathbf{p}$ implicit in Eq.(26) for the interference function $I(\mathbf{p}, \mathbf{b})$, have appeared as oscillations in space $\mathbf{r}$ for a “one frame snap shot” [9] of $\rho_0(\mathbf{r}, t_{\text{fixed}})$ fixed in time. Interference is also present as a time oscillation in $\rho_0(\mathbf{r}_{\text{fixed}}, t)$ for a movie picture of atoms in the neighborhood of a fixed point in space[16].

The point is that the momentum, position and time are related by the “classical path rule” $\mathbf{r} = \mathbf{v}t = (\mathbf{p}t/M)$, but with a *weighted momentum distribution*. This *essence* of the diffraction of particle beams as described by quantum mechanical amplitude superposition, should also be possible to observe in systems for which the one rotates the potential about an axis displaced from the symmetry axis. Mesoscopic quantum interference should also be present in TOP traps[7].

REFERENCES

1. I. M. Khalatnikov, *Introduction to the Theory of Superfluidity*, W. A. Benjamin Inc., New York (1965).
2. L. D. Landau, *J. Phys.* **5**, 71 (1941).
3. N. N. Bogoluibov, *J. Phys.* **11**, 23 (1947).
4. S. Giorgini, L. P. Pitaevskii and S. Stringari, *J. Low Temp. Phys.* **109**, 309 (1997).
5. M. Narasahewski, H. Wallis and A. Schenzle *Phys. Rev. A* **54** 2185 (1996).
6. W. Ketterle, M. R. Andrews, K. B. Davis, D. S. Durfee, D. M. Kurn, M. -O. Mewes and N. J. van Druten, *Physica Scripta* **T66**, 31 (1996).
7. M. H. Anderson, J. R. Ensher, M. R. Matthews, C. E. Wieman and E. A. Cornell, *Science* **269**, 198 (1995);
8. C. C. Bradley, C. A. Sackett, J. J. Tollet and R. G. Hulet, *Phys. Rev. Lett.* **75** 1687 (1995).
9. M. R. Andrews, C. G. Townsend, H. -J. Miesner, D. S. Durfee, D. M. Kurn and W. Ketterle, *Science* **275**, 637 (1997).
10. A. S. Davydov, *Quantum Mechanics*, Sec. 92, pp. 394-96, Pergamon Press, Oxford (1985).
11. R. P. Feynman, *Rev. Mod. Phys.*, **20**, 267 (1948).
12. K. B. Davis, M. -O. Mewes, M. R. Andrews, N. J. van Druten, D. S. Durfee, D. M. Kurn and W. Ketterle, *Phys. Rev. Lett.* **75**, 3969 (1995).
13. A. Tonomura, N. Osakabe, T. Matsuda, T. Kawasaki, J. Endo, S. Yano and H. Yamada, *Phys. Rev. Lett.* **56**, 792 (1986).
14. E. P. Gross, *Il Nouvo Cimento* **20**, 454 (1961).
15. L. P. Pitaevskii, *JETP* **13**, 451 (1961).
16. D. M. Stamper-Kurn, H.-J. Miesner, S. Inouye, M. R. Andrews and W. Ketterle, cond-mat/9801262 (1998).